

PERTURBATIVE EVALUATION OF THE ZERO-POINT FUNCTION FOR SELF-INTERACTING SCALAR FIELD ON A MANIFOLD WITH BOUNDARY

GEORGE TSOUPROS
SCHOOL OF PHYSICS
THE UNIVERSITY OF NEW SOUTH WALES
NSW 2052
AUSTRALIA

ABSTRACT. The character of quantum corrections to the gravitational action of a conformally invariant field theory for a self-interacting scalar field on a manifold with boundary is considered at third loop-order in the perturbative expansion of the zero-point function. Diagrammatic evaluations and higher loop-order renormalisation can be best accomplished on a Riemannian manifold of positive constant curvature accommodating a boundary of constant extrinsic curvature. The associated spherical formulation for diagrammatic evaluations reveals a non-trivial effect which the topology of the manifold has on the vacuum processes and which ultimately dissociates the dynamical behaviour of the quantised field from its behaviour in the absence of a boundary. The first surface divergence is evaluated and the necessity for simultaneous renormalisation of volume and surface divergences is shown.

I. Introduction

The development of Euclidean Quantum Gravity has emphasised the importance of any investigation relevant to the dynamical behaviour of quantised matter fields on a manifold with boundary by revealing the latter's physical significance in the quantum cosmological context of semiclassical tunnelling. Specifically, instanton-related considerations in quantum and inflationary cosmology lend particular importance to the issue of radiative contributions to a semiclassical tunnelling geometry of positive constant curvature effected by quantised matter. The presence of a boundary drastically alters the dynamical behaviour of the quantum field. In addition to the contributions which any divergence stemming from vacuum effects at any loop-order receives from the Riemann curvature of the background geometry the same divergence receives contributions from the extrinsic curvature of the boundary. This effect which has been studied at one-loop order through heat kernel techniques [1] will be shown in this project to be relevant to any loop-order. Its occurrence is exemplary of the potential effect which the topology of a manifold has on the short distance dynamical behaviour of the quantum field defined on it.

1991 *Mathematics Subject Classification.* 49Q99, 81T15, 81T18, 81T20.
present e-mail address: gts@phys.unsw.edu.au.

Any evaluation of higher loop-order vacuum effects generated on the dynamical behaviour of quantised matter on a manifold with boundary necessitates diagrammatic techniques. Diagrammatic evaluations and higher loop-order renormalisation can be best accomplished on a Riemannian manifold of positive constant curvature accommodating a boundary of constant extrinsic curvature. The advantage of working on a Euclidean n -dimensional spherical cap C_n stems from its underlying symmetry which allows for the exploitation of the method of images, thereby allowing for the exploitation of the emerging spherical symmetry in the evaluation of radiative effects. In the physically important case of a massless scalar field conformally coupled to the background geometry of C_n with the Dirichlet condition specified on the boundary ∂C_n such an approach has resulted in an exact expression for the massless propagator [2] [3]. This, in turn, allows for the development of the diagrammatic techniques necessary for the evaluation of higher loop-order radiative contributions to the associated effective action for a self-interacting conformal scalar field at $n = 4$. A prerequisite for such an evaluation is a spherical formulation of the Feynman rules on C_n . Such a formulation has been advanced in the context of the method of images by exploiting the association of the dynamical behaviour of the conformal scalar field on C_n to that of the same field on the Euclidean n -sphere S_n [3]. Consequent upon such a formulation was, in addition, a confirmation of the results obtained in [2] to the effect that, at any loop-order, no independent renormalisation is necessary. The volume-related terms in the semiclassical action receive infinite contributions from vacuum effects simultaneously with the boundary-related terms. In this respect, the stated spherical formulation confirmed, in addition, the potential for non-trivial radiative generation of surface counterterms in the effective action.

The assessment of the stated boundary-related and vacuum effects will be pursued, in what follows, by exploiting the results obtained in [3] in order to evaluate the zero-point function at third loop-order (second order in the self-coupling) for a self-interacting massless scalar field conformally coupled to the background geometry of C_4 . Such an evaluation will also constitute the incipient point of a renormalisation program to the same order. From the outset, all divergences in the gravitational action generated by scalar loops on a four dimensional manifold are, on dimensional grounds, expected to be proportional to the cosmological constant Λ , to the Ricci scalar R , to R^2 , and to the Ricci and Riemann tensor-related contractions $R_{\mu\nu}R^{\mu\nu}$ and $R_{\alpha\beta\gamma\delta}R^{\alpha\beta\gamma\delta}$ respectively. Suitable linear combinations of these terms are also possible [4]. If a boundary is present additional divergent contributions involving the extrinsic curvature K of that boundary are expected [2]. The advantage of evaluating such gravitational divergences on a manifold of constant curvature is that the concomitant spherical formulation of the theory determines each Green function as an exact function of the constant Ricci scalar R [5]. This result has both formal and perturbative significance and constitutes, for that matter, the primary motivation for the entire spherical formulation of euclidean quantum field theory [6], [7], [8], [9], [10]. Effectively, only volume-related counterterms proportional to Λ , R , R^2 and RK^2 are allowed on C_4 . This is the case because, perturbatively, the only source of volume divergences are terms in the relevant expansions of the Green functions.

The effect which a boundary in space-time has on the dynamical behaviour of quantised matter can be best appreciated through a comparison with the same dynamical behaviour in the absence of a boundary. In the present case, such a comparison is facilitated by the fact that both C_4 and S_4 are manifolds of positive constant curvature. In what follows it will be shown that the divergences inherent in the zero-point function evaluated at three-loop level receive a non-trivial contribution from the boundary ∂C_4 which, effectively, dissociates the dynamical behaviour of the quantised scalar field on C_4 from that on S_4 . It is also worth noting in this respect, that since divergences in the evaluation of loop effects stem essentially from the coincidence of integration points in the relevant Feynman diagrams, the emergence of divergences proportional to K signifies a non-trivial effect which the topology of a bounded manifold has on the cut-off scale behaviour of the quantised matter fields. The announced simultaneous renormalisation of volume and surface terms will, also, become manifest in the context of the perturbative calculation pursued in this project. Moreover, the stated potential of the theory for non-trivial generation of surface counterterms in the effective action will be realised in the evaluation of the first surface divergence on a manifold with boundary.

II. The Zero Point Function

The unique conformally invariant scalar action on a n -dimensional Riemannian manifold of positive constant embedding radius a is [5]

$$(1) \quad S[\Phi] = \int_C d\sigma \left[\frac{1}{2} \frac{1}{2a^2} \Phi (L^2 - \frac{1}{2} n(n-2)) \Phi - \frac{\lambda}{\Gamma(p+1)} \Phi^p \right]$$

provided that $p = \frac{2n}{n-2}$, $n > 2$. $L_{\mu\nu}$ is the generator of rotations

$$(2) \quad L_{\mu\nu} = \eta_\mu \frac{\partial}{\partial \eta_\nu} - \eta_\nu \frac{\partial}{\partial \eta_\mu}$$

on the relevant embedded manifold. In the physically relevant case of $n = 4$ the self-coupling is that of Φ^4 . Any n -dimensional Riemannian manifold of positive constant embedding radius a represents, through the usual rotation in imaginary time, the Euclidean version of the associated segment of de Sitter space. On any such manifold the Ricci scalar R admits the constant value [5]

$$(3) \quad R = \frac{n(n-1)}{a^2}$$

In conformity with the considerations hitherto outlined the Riemannian manifold relevant to (1) will be specified to be that of a spherical n -cap C_n considered as a manifold of positive constant curvature embedded in a $(n+1)$ -dimensional Euclidean space and bounded by a $(n-1)$ -sphere of positive constant extrinsic curvature K (diverging normals). In effect, C_n is characterised by spherical $(n-1)$ -dimensional sections of constant Euclidean time τ . The Einstein-Hilbert action, being the gravitational component of the conformally invariant (semi)classical action, necessitates an additional term of the form

$\int_{\partial C} K \Phi^2$ enforced by the presence of the boundary ∂C [2]. As a result, the Einstein-Hilbert action at $n = 4$ assumes the form

$$(4) \quad S_{EH} = -\frac{1}{16\pi G} \int_C d\sigma (R - 2\Lambda) + \int_{\partial C} d^3x \sqrt{h} K \Phi^2$$

with $d\sigma = a^n d\Omega_{n+1}$ being the element of surface area of the n -sphere embedded in $n + 1$ dimensions and with the three-dimensional boundary hypersurface on which the induced metric is h_{ij} being characterised by an extrinsic curvature $K_{ij} = \frac{1}{2}(\nabla_i n_j + \nabla_j n_i)$ the trace of which is $K = h^{ij} K_{ij} > 0$. Although the surface term in (4) stems from a variational demand posed on S_{EH} it is not purely geometrical. For that matter it is considered an additional sector to the scalar component of the action expressed in (1). The action $S[\Phi]$ for the specified theory is the sum-total of all sectors in (1) and (4). In addition to the cosmological constant and scalar curvature featured in S_{EH} the semiclassical approach, which differentiates between the matter and gravitational degrees of freedom by quantising the former on a fixed geometrical background, is known to result in all possible volume divergences which are dimensionally consistent in the gravitational action. These higher-order corrections are effected by vacuum processes of the interacting field [13]. The immediate issue which the presence of a boundary raises perturbatively is the possibility of additional divergent contributions in the bare gravitational action featuring the extrinsic curvature K . In fact, such contributions to the two-point function - and, for that matter, to the bare scalar action - are expected past one-loop order [2].

All Green functions of Φ^4 theory on C_4 are generated by the Euclidean functional integral

$$(5) \quad Z[J] = \int D[\Phi] e^{S[\Phi] + J\Phi}$$

through functional differentiation at $J = 0$. This generating functional is the mathematical expression for the transition amplitude between vacuum states defined on C_4 [4]. The disconnected components of this generating functional in curved space-times - represented by the zero-point function - invariably have non-trivial contributions to that transition amplitude and, for that matter, to the effective action. This is, essentially, due to graviton contributions which the matter propagators receive from the semiclassical background geometry and which, for that matter, render a purely geometric character to any divergences inherent in the zero-point function. Although stemming from the dynamical behaviour of the matter fields these divergences result in the infinite redefinitions in the gravitational component of the semiclassical action.

Pursuant to the spherical formulation relevant to diagrammatic evaluations on C_n developed in [3] all calculations will be advanced in configuration space. The first two terms in the perturbative expansion of the zero-point function in powers of $\hbar$ (loop expansion) are diagrammatically represented by the graphs of fig.(1). The “bubble” diagrams in fig.(1a) and fig.(1b) account for the one-loop contribution to the zero-point function of the theory. Their simultaneous presence in any curved space-time is expected on the basis of general theoretical considerations [4]. They are characterised by the absence of interaction vertices and, on power counting grounds, are responsible for the simultaneous

one-loop contributions to volume and boundary effective Einstein-Hilbert action on any manifold with boundary [16]. They have been shown to be finite provided that dimensional regularisation is used [2]. The latter technique regulates all ultraviolet divergences through an analytical extension of the space-time dimensionality to an arbitrary value n [11].




FIGURE 1. $O(\hbar^3)$ -related contributions to the zero-point function

The presence of the self-coupling in the semi-classical scalar action generates additional contributions to the zero-point function at two and three loop-level represented by fig.(1d) and fig.1(f). They are finite in the context of dimensional regularisation and, for that matter, just as trivial as those relevant to fig.(1a) and fig.(1b)




FIGURE 2. $O(\hbar^3)$ -related finite contributions to the zero-point function

The remaining diagram in fig.(1c) represents the first potentially non-trivial contribution in the perturbative expansion of the zero-point function. It is of order $\hbar^3$ and features two interaction vertices. In view of the minimal subtraction-related expansion

$$(6) \quad \lambda_0 = \mu^{4-n} \left[\lambda + \sum_{\nu=1}^{\infty} \frac{a_{\nu}(\lambda)}{(4-n)^{\nu}} \right]$$

which the bare self-coupling λ_0 admits in powers of the renormalised self-coupling λ [11] the multiplicative factor of $(-\lambda_0)$ dictated by the Feynman rules for each of the two vertices of fig.(1c) amounts, at that three-loop order, to λ^2 [5]. In effect, the diagram in question featuring four internal propagators joining the vertices η and η' results in the following mathematical expression for the stated $O(\hbar^3)$ -term

$$(7) \quad I_c = \lambda^2 B_n^4 \int d^n \eta d^n \eta' [D_c^{(n)}(\eta, \eta')]^4$$

with

$$(8) \quad B_n = \frac{\Gamma(\frac{n}{2} - 1)}{4\pi^{\frac{n}{2}}}$$

The multiplicative factor of B_n is the symmetry factor associated with the propagator $D_c^{(n)}(\eta, \eta')$ for the conformal field Φ on C_n which was shown to be [2], [3]

$$(9) \quad D_c^{(n)}(\eta, \eta') = \frac{\Gamma(\frac{n}{2} - 1)}{4\pi^{\frac{n}{2}}} \frac{1}{|\eta - \eta'|^{n-2}} - \frac{\Gamma(\frac{n}{2} - 1)}{4\pi^{\frac{n}{2}}} \frac{1}{|\frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta'|^{n-2}}$$

with a_B and $a_{\eta'}$ being the geodesic distances between the cap's pole and its boundary as well as between the former and point η' respectively. Replacing this expression in that for I_c and expanding yields

$$(10) \quad \begin{aligned} I_c = \lambda^2 B_n^4 \int_C d^n \eta d^n \eta' [& |\eta - \eta'|^{4(2-n)} + 4|\eta - \eta'|^{3(2-n)} |\frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta'|^{2-n} + \\ & 6|\eta - \eta'|^{2(2-n)} |\frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta'|^{2(2-n)} + \\ & 4|\eta - \eta'|^{2-n} |\frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta'|^{3(2-n)} + |\frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta'|^{4(2-n)}] \end{aligned}$$

All five terms in this double volume integral will have to be examined separately. Their evaluation necessitates the results obtained through a spherical formulation for the mathematical expressions of Feynman diagrams on C_n [3]. In the ensuing calculations use will be made of the expressions

$$(11) \quad [(\eta - \eta')^2]^\nu = \sum_{N=0}^{\infty} \sum_{\alpha=0}^N \frac{(2a)^{2\nu+n} \pi^{\frac{n}{2}} \Gamma(\nu + \frac{n}{2}) \Gamma(N - \nu)}{\Gamma(N + n + \nu) \Gamma(-\nu)} Y_\alpha^N(\eta) Y_\alpha^N(\eta')$$

$$(12) \quad [|\frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta'|^2]^\nu = \sum_{N=0}^{N_0} \sum_{\alpha=0}^N \frac{(2a)^{2\nu+n} \pi^{\frac{n}{2}} \Gamma(\nu + \frac{n}{2} + \frac{1}{N_0}) \Gamma(N - \nu + \frac{1}{N_0})}{\Gamma(N + n + \nu + \frac{1}{N_0}) \Gamma(-\nu)} Y_\alpha^N(\eta) Y_\alpha^N(\eta')$$

for the powers appearing in (10). Moreover, in the context of the spherical formulation developed in [3] the orthonormality condition for the n-dimensional spherical harmonics defined on S_n

$$(13) \quad \int_S d^n \eta Y_\alpha^N(\eta) Y_{\alpha'}^{N'}(\eta) = \delta_{NN'} \delta_{\alpha\alpha'}$$

is, necessarily, replaced on C_n in terms of Gegenbauer polynomials $C_m^p(\cos\theta)$ with their explicit dependence on the angles associated with the embedding $(n+1)$ -vector η

$$\begin{aligned} \int_C d^n \eta Y_\alpha^N(\eta) Y_{\alpha'}^{N'}(\eta) &= AK a (\sin\theta_n^0)^{m_1+l_1} C_{N-m_1}^{m_1+\frac{n-1}{2}}(\cos\theta_n^0) C_{N'-l_1}^{l_1+\frac{n-1}{2}}(\cos\theta_n^0) \delta^{m_1 l_1} \delta_{aa'} + \\ &AB \prod_{k=1}^{n-2} \int_0^\pi C_{m_k-m_{k+1}}^{m_{k+1}+\frac{n-2}{2}}(\cos\theta_{n-k}) C_{l_k-l_{k+1}}^{l_{k+1}+\frac{n-2}{2}}(\cos\theta_{n-k}) [\sin\theta_{n-k}]^{m_{k+1}+l_{k+1}+n-3} d\theta_{n-k} \times \\ &\int_0^{2\pi} \frac{e^{i(\pm m_{n-1} \mp l_{n-1})\theta_1}}{\sin\theta_1} \end{aligned} \quad (14a)$$

with θ_n^0 being the angle associated with ∂C_n , with

$$m_1 = N \cos\theta_n^0; \quad l_1 = N' \cos\theta_n^0 \quad (14b)$$

being the degrees of spherical harmonics defined on ∂C_n , with

$$\delta_{\alpha\alpha'} = \delta^{m_1 l_1} \delta_{aa'} \quad (14c)$$

and with

$$A = \frac{1}{(N' + \frac{n}{2} - 1)(N' + \frac{n}{2}) - (N + \frac{n}{2} - 1)(N + \frac{n}{2})}; \quad N \neq N' \quad (14d)$$

and

$$\begin{aligned} B &= 2 [m_1 (\sin\theta_n^0)^{m_1+l_1+n-2} (\cos\theta_n^0) C_{N-m_1}^{m_1+\frac{n-1}{2}}(\cos\theta_n^0) - \\ &(\sin\theta_n^0)^{m_1+l_1+n} (2m_1+n-1) C_{N-m_1-1}^{m_1+\frac{n-1}{2}+1}(\cos\theta_n^0)] C_{N'-l_1}^{l_1+\frac{n-1}{2}}(\cos\theta_n^0) \end{aligned} \quad (14e)$$

The first term in the integrand of (10) stems exclusively from the fundamental part of the propagator $D_c^{(n)}(\eta, \eta')$ and is, for that matter, identical to the corresponding term which upon integration over the volume of S_n yields the entire contribution to the zero-point function at $O(\lambda^2)$ on the Euclidean de Sitter space. As, in the present case, the integral is taken over a bounded manifold a comparison of this mathematical expression on S_4 and C_4 would be explicitly indicative of the mathematical origin of boundary-related contributions to divergences stemming from vacuum effects. The corresponding integral for the contribution to the zero-point function on S_n , for that matter, is

$$I_S = \lambda^2 B_n^4 \int_S d^n \eta d^n \eta' [|\eta - \eta'|^2]^{\frac{8-4n}{2}} \quad (15)$$

which, in the context of (11), yields

$$(16) \quad I_S = \lambda^2 B_n^4 \sum_{N=0}^{\infty} \sum_{\alpha=0}^N \frac{(2a)^{8-3n} \pi^{\frac{n}{2}} \Gamma(\frac{8-3n}{2}) \Gamma(N - \frac{8-4n}{2})}{\Gamma(N + \frac{8-2n}{2}) \Gamma(\frac{4n-8}{2})} \int_S d^n \eta d^n \eta' Y_\alpha^N(\eta) Y_\alpha^N(\eta')$$

whereupon, due to the orthonormality condition (13) and the fact that Y_0^0 relates to the embedding radius a and to the solid angle Ω_{n+1} for S_n through [12]

$$(17) \quad Y_0^0 = \frac{1}{\sqrt{a^n \Omega_{n+1}}}$$

it further reduces to

$$I_S = \lambda^2 2^{-3n} \pi^{-\frac{3n}{2}} a^{8-3n} \frac{[\Gamma(\frac{n}{2} - 1)]^4 \Gamma(4 - \frac{3n}{2})}{\Gamma(4 - n)} \int_S d^n \eta$$

where use has been made of (9) as well. It is evident that at $n \rightarrow 4$ the pole contained in $\Gamma(4 - \frac{3n}{2})$ cancels against that in $\Gamma(4 - n)$. Setting $\epsilon = 4 - n$ and using the standard expansion

$$(18) \quad \Gamma(\epsilon) = \frac{1}{\epsilon} - \gamma + \frac{\frac{\pi^2}{6} + \gamma^2}{2} \epsilon + \dots$$

for $\epsilon \ll 1$ the final expression at $n \rightarrow 4$ is

$$(19) \quad I_S = \lambda^2 \frac{1}{3^3 2^{16} \pi^6} R^2$$

where use has been made of (3) and a multiplicative factor amounting to the volume of S_n is implied.

The result expressed in (19) amounts to a finite contribution to the gravitational effective action on S_4 . It becomes evident, for that matter, that the zero-point function has no divergent contribution to the gravitational component of the action on S_4 to second order in the self-coupling [$O(\hbar^3)$]. This result is consistent with [13] where, in fact, it was shown that on a general four-dimensional unbounded curved manifold vacuum effects due to a ϕ^4 self-interaction do not generate divergent contributions to the R^2 sector of the gravitational action before the fifth order in λ , provided that dimensional regularisation is used.

The same procedure applied on the same term integrated over the volume of C_n results in the following expression for the first term in (10)

$$(20) \quad \lambda^2 B_n^4 \int_C d^n \eta d^n \eta' [|\eta - \eta'|^2]^{\frac{8-4n}{2}} = \lambda^2 B_n^4 \sum_{N=0}^{\infty} \sum_{\alpha=0}^N \frac{(2a)^{8-3n} \pi^{\frac{n}{2}} \Gamma(\frac{8-3n}{2}) \Gamma(N - \frac{8-4n}{2})}{\Gamma(N + \frac{8-2n}{2}) \Gamma(\frac{4n-8}{2})} \int_C d^n \eta d^n \eta' Y_\alpha^N(\eta) Y_\alpha^N(\eta')$$

where, again, the double integral is to be tackled on the basis of (17)

$$\int_C d^n \eta Y_\alpha^N(\eta) = \sqrt{a^n \Omega_{n+1}} \int_C d^n \eta Y_\alpha^N(\eta) Y_0^0$$

on the understanding that the orthonormality condition (13) on S_n is replaced by the condition (14) on C_n . In this case, however, the constancy of Y_0^0 enforces a vanishing result on the intractable second term of (14a). Rather than deal with that term itself a faster expedient to this result is provided directly by the reduction formula [3]

$$(21) \quad \int_C d^n \eta Y_\alpha^N(\eta) Y_{\alpha'}^{N'}(\eta) = Aa^2 \oint_{\partial C} d^{n-1} \eta [KY_\alpha^N(\eta) Y_{\alpha'}^{N'}(\eta) + 2n_p Y_{\alpha'}^{N'}(\eta) D_p Y_\alpha^N(\eta)]$$

which is, itself, a condition for (14) and in whose context it becomes obvious that for $Y_{\alpha'}^{N'} = Y_0^0$ the stated second term merely contributes a factor of $-2KY_\alpha^N(\eta)Y_0^0$ in the integrand. Consequently,

$$\int_C d^n \eta Y_\alpha^N(\eta) Y_0^0 = -aK \frac{(\sin \theta_n^0)^{m_1} C_{N-m_1}^{m_1 + \frac{n-1}{2}} (\cos \theta_n^0) C_0^{\frac{n-1}{2}}}{(\frac{n}{2} - 1)\frac{n}{2} - (N + \frac{n}{2} - 1)(N + \frac{n}{2})} \delta^{m_1 0} \delta_{a0}; \quad N \neq 0$$

which, through the relation [12]

$$C_0^k(x) = 1; k > -\frac{1}{2}$$

reduces to

$$\int_C d^n \eta Y_\alpha^N(\eta) Y_0^0 = -aK \frac{1}{(\frac{n}{2} - 1)\frac{n}{2} - (N + \frac{n}{2} - 1)(N + \frac{n}{2})} C_N^{\frac{n-1}{2}} (\cos \theta_n^0)$$

so that the double integral over C_n in (20) eventually becomes

$$(22) \quad \int_C d^n \eta d^n \eta' Y_\alpha^N(\eta) Y_\alpha^N(\eta') = \frac{a^2 K^2}{[(\frac{n}{2} - 1)\frac{n}{2} - (N + \frac{n}{2} - 1)(N + \frac{n}{2})]^2} [C_N^{\frac{n-1}{2}} (\cos \theta_n^0)]^2 a^n \Omega_{n+1}$$

In the context of (20), (22), (14c) and (8) the first term in (10) assumes the form

$$(23) \quad \lambda^2 B_n^4 \int_C d^n \eta d^n \eta' [|\eta - \eta'|]^{\frac{8-4n}{2}} = \lambda^2 a^{10-2n} K^2 \Omega_{n+1} \left[\frac{\Gamma(\frac{n}{2} - 1)}{4\pi^{\frac{n}{2}}} \right]^4 \sum_{N=0}^{\infty} \frac{[C_N^{\frac{n-1}{2}} (\cos \theta_n^0)]^2}{[(\frac{n}{2} - 1)\frac{n}{2} - (N + \frac{n}{2} - 1)(N + \frac{n}{2})]^2} \frac{(2)^{8-3n} \pi^{\frac{n}{2}} \Gamma(\frac{8-3n}{2}) \Gamma(N - \frac{8-4n}{2})}{\Gamma(N + \frac{8-2n}{2}) \Gamma(\frac{4n-8}{2})}; \quad N \neq 0$$

whereupon setting $\epsilon = 4 - n$ and using (18) prior to taking $\epsilon \rightarrow 0$ renders the first term of (10) in the form

$$\lambda^2 B_n^4 \int_C d^n \eta d^n \eta' [|\eta - \eta'|]^{4(2-n)} \rightarrow$$

$$(24) \quad \lambda^2 a^2 K^2 \Omega_5 2^{-13} 3^{-2} \pi^{-6} \left[\sum_{N=1}^{\infty} [C_N^{\frac{3}{2}}(\cos\theta_4^0)]^2 \frac{(N+1)(N+2)}{N(N+3)} \right] \frac{1}{\epsilon} + F.T.$$

where use has, also, been made of the standard expressions for Γ functions of an integer argument m and of a complex argument z respectively

$$\Gamma(m) = (m-1)! ; \quad z\Gamma(z) = \Gamma(z+1)$$

The series of any power of $C_N^k(x)$ over the degree N for a fixed order k has a radius of convergence between $(-1, 1)$ [12]. For that matter, the series featured in (24), is convergent and the first term of (10) displays, as a consequence, a simple pole at $n \rightarrow 4$.

The physical interpretation of (24) becomes obvious through the observation that the volume of the embedded S_4 relates to the solid angle Ω_5 through

$$\int_C d^4\eta + \int_{S-C} d^4\eta = a^4 \Omega_5$$

with

$$\int_{S-C} d^4\eta = V_c \int_C d^4\eta; \quad V_c > 0$$

which renders (24) exclusively in terms of volume integration over C_4

$$\lambda^2 B_n^4 \int_C d^n\eta d^n\eta' [|\eta - \eta'|]^{4(2-n)} \rightarrow$$

$$\lambda^2 \frac{1}{a^2} K^2 2^{-13} 3^{-2} \pi^{-6} \left[\sum_{N=1}^{\infty} [C_N^{\frac{3}{2}}(\cos\theta_4^0)]^2 \frac{(N+1)(N+2)}{N(N+3)} \right] \frac{1}{\epsilon} \int_C d^4\eta + F.T.$$

allowing for a multiplicative factor which depends exclusively on the volume of C_4 . Use of (3) then finally yields at $n \rightarrow 4$

$$\lambda^2 B_n^4 \int_C d^n\eta d^n\eta' [|\eta - \eta'|]^{4(2-n)} \rightarrow$$

$$(25) \quad \lambda^2 \frac{1}{2^{15} 3^3 \pi^6} \left[\sum_{N=1}^{\infty} [C_N^{\frac{3}{2}}(\cos\theta_4^0)]^2 \frac{(N+1)(N+2)}{N(N+3)} \right] (RK^2) \frac{1}{\epsilon} \int_C d^4\eta + F.T.$$

In sharp contrast with (19) this result reveals that volume integration over C_n of the fourth power of the fundamental part of the propagator in (9) entails a pole at the limit of space-time dimensionality. The absence of any divergent contribution to the bare gravitational action on S_4 at the same loop-order reveals that the underlying source of the pole revealed at $n \rightarrow 4$ is ∂C_4 , an effect which is mathematically reflected in the explicit presence of the boundary's extrinsic curvature K in (25). In [3] the case was made to the effect that the reduction of the complete orthonormality condition on S_n expressed by (13) to the partial orthonormality condition on C_n expressed by (14) is responsible for the contributions which any divergence on S_4 receives due to the

presence of ∂C_4 . A comparison of the calculational context of (19) with that of (25) reveals indeed that, in the context of dimensional regularisation, the finite contribution to the effective action on S_4 emerges as a result of the pole cancellation relevant to the numerator-related $\Gamma(\frac{8-3n}{2})$ and the denominator-related $\Gamma(N + \frac{8-2n}{2})$ in (16). That cancellation, however, which essentially stems from (13) on S_4 is unattainable on C_4 where that orthonormality condition “breaks down” allowing, as a result, for an infinite number of N -dependent terms in addition to the cancellation-associated $N = 0$ term. Moreover, since the replacement of (13) by (14) responsible for that effect is, itself, a direct consequence of the presence of the boundary ∂C_4 it becomes evident that all boundary-related contributions to divergences already present on S_4 are, necessarily, effected by the volume integral which, in the context of the Feynman rules, stems from vertex integration in the configuration space of C_4 [3].

The second, third and fourth term in (10) involve products of powers associated with both the fundamental and boundary part of the propagator in (9). They necessitate, for that matter, a simultaneous use of (11) and (12) on the understanding that at the dimensional limit $n \rightarrow 4$ all divergences stem exclusively from (11), a fact which is ensured by the presence of the inverse of the upper limit N_0 related to the cut-off angular momentum for image propagation in the arguments of the Γ functions in (12) [3]. The simultaneous substitutions through (11) and (12) are expected to result in mathematical expressions which are substantially more involved than that of the, hitherto tackled, first term. These expressions necessarily relate to the residues of the resulting poles and to finite terms of the associated expansions and do not, for that matter, complicate the pole-structure of the theory. In effect, a direct application of (11), (12) and (8) on the second term of (10) results in

$$\begin{aligned}
 & \lambda^2 B_n^4 4 \int_C d^n \eta d^n \eta' |\eta - \eta'|^{3(2-n)} \left| \frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta' \right|^{2-n} = \lambda^2 \left[\Gamma\left(\frac{n}{2} - 1\right) \right]^4 \pi^{-2n} 2^{-6} \times \\
 & \sum_{N=0}^{\infty} \sum_{\alpha=0}^N \frac{(2a)^{6-2n} \pi^{\frac{n}{2}} \Gamma\left(\frac{6-2n}{2}\right) \Gamma\left(N - \frac{6-3n}{2}\right)}{\Gamma\left(N + \frac{6-n}{2}\right) \Gamma\left(\frac{3n-6}{2}\right)} \sum_{N'=0}^{N_0'} \sum_{\alpha'=0}^{N'} \frac{(2a)^2 \pi^{\frac{n}{2}} \Gamma\left(1 + \frac{1}{N_0'}\right) \Gamma\left(N' - \frac{2-n}{2} + \frac{1}{N_0'}\right)}{\Gamma\left(N' + \frac{2+n}{2} + \frac{1}{N_0'}\right) \Gamma\left(\frac{n-2}{2}\right)} \times \\
 & \int_C d^n \eta Y_{\alpha}^N(\eta) Y_{\alpha'}^{N'}(\eta) \int_C d^n \eta' Y_{\alpha}^N(\eta') Y_{\alpha'}^{N'}(\eta')
 \end{aligned}
 \tag{26}$$

Since each of the integrals over the product of spherical harmonics is directly expressed by (14) it is

$$\begin{aligned}
 & \int_C d^n \eta Y_{\alpha}^N(\eta) Y_{\alpha'}^{N'}(\eta) \int_C d^n \eta' Y_{\alpha}^N(\eta') Y_{\alpha'}^{N'}(\eta') = \\
 & \frac{1}{[(N' + \frac{n}{2} - 1)(N' + \frac{n}{2}) - (N + \frac{n}{2} - 1)(N + \frac{n}{2})]^2} \times \\
 & [a^2 K^2 F^2(n) \delta_{\alpha\alpha'} + 2aK[F(n)B(n)H(n)]\delta_{\alpha\alpha'} + [B(n)H(n)]^2]
 \end{aligned}
 \tag{27}$$

with the condition $N \neq N'$ and with

$$F(n) = (\sin\theta_n^0)^{2m_1} C_{N-m_1}^{m_1+\frac{n-1}{2}}(\cos\theta_n^0) C_{N'-m_1}^{m_1+\frac{n-1}{2}}(\cos\theta_n^0) ; m_1 = N\cos\theta_n^0, l_1 = N'\cos\theta_n^0$$

and

$$H(n) = \prod_{k=1}^{n-2} \int_0^\pi C_{m_k-m_{k+1}}^{m_{k+1}+\frac{n-2}{2}}(\cos\theta_{n-k}) C_{l_k-l_{k+1}}^{l_{k+1}+\frac{n-2}{2}}(\cos\theta_{n-k}) [\sin\theta_{n-k}]^{m_{k+1}+l_{k+1}+n-3} d\theta_{n-k} \times$$

$$\int_0^{2\pi} \frac{e^{i(\pm m_{n-1} \mp l_{n-1})\theta_1}}{\sin\theta_1}$$

Substituting that expression in (26) and taking the limit $4-n=\epsilon \rightarrow 0$ results in a simple pole contained in $\Gamma(\frac{6-2n}{2})$

$$\lambda^2 B_n^4 4 \int_C d^n \eta d^n \eta' |\eta - \eta'|^{3(2-n)} \left| \frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta' \right|^{2-n} \rightarrow$$

$$\lambda^2 \frac{(-1)}{2^7 \pi^4} \sum_{N=0}^{\infty} \sum_{N'=0}^{N'_0} \frac{\Gamma(N+3)}{\Gamma(N+1)} \frac{\Gamma(1+\frac{1}{N'_0}) \Gamma(N'+1+\frac{1}{N'_0})}{\Gamma(N'+3+\frac{1}{N'_0})} \frac{1}{[N'^2 - N^2 + 3(N' - N)]^2} \times$$

$$\left[a^2 K^2 F^2(4) \sum_{\alpha=0}^N \sum_{\alpha'=0}^{N'} \delta_{\alpha\alpha'} + 2aK[F(4)B(4)H(4)] \sum_{\alpha=0}^N \sum_{\alpha'=0}^{N'} \delta_{\alpha\alpha'} + [B(4)H(4)]^2 \right] \frac{1}{\epsilon} + F.T.$$

which, in view of the double summation over the Kronecker symbol and the condition $N \neq N'$, amounts to

$$\lambda^2 B_n^4 4 \int_C d^n \eta d^n \eta' |\eta - \eta'|^{3(2-n)} \left| \frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta' \right|^{2-n} \rightarrow$$

$$\lambda^2 \frac{(-1)}{2^7 \pi^4} \sum_{N=0}^{\infty} \sum_{N'=0}^{N'_0} \frac{\Gamma(N+3)}{\Gamma(N+1)} \frac{\Gamma(1+\frac{1}{N'_0}) \Gamma(N'+1+\frac{1}{N'_0})}{\Gamma(N'+3+\frac{1}{N'_0})} \frac{1}{[N'^2 - N^2 + 3(N' - N)]^2} \times$$

$$[a^2 K^2 F^2(4) N' + 2aK[F(4)B(4)H(4)] N' + [B(4)H(4)]^2] \frac{1}{\epsilon} + F.T.$$

whereupon through use of

$$\int_C d^4 \eta = a^4 \Omega_5^{(C)}$$

in the first and third term of the geometry-dependent multiplicative factor and

$$\oint_{\partial C} d^3 \eta = (a \sin(\theta_4^0))^3 \Omega_4$$

in the second term of that factor and through (3) for $n=4$ the result for the second term in question is

$$\begin{aligned}
& \lambda^2 B_n^4 4 \int_C d^n \eta d^n \eta' |\eta - \eta'|^{3(2-n)} \left| \frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta' \right|^{2-n} \rightarrow \\
& \lambda^2 \frac{(-1)}{2^7 \pi^4} \sum_{N=0}^{\infty} \sum_{N'=0}^{N'_0} \frac{\Gamma(1 + \frac{1}{N'_0}) \Gamma(N' + 1 + \frac{1}{N'_0})}{\Gamma(N' + 3 + \frac{1}{N'_0})} \frac{(N+1)(N+2)}{[N'^2 - N^2 + 3(N' - N)]^2} \times \\
& \left[\frac{1}{3} \frac{1}{2^2} F^2(4) N'(RK^2) (\Omega_5^{(C)})^{-1} \int_C d^4 \eta + \frac{1}{3} \frac{1}{2} [F(4) B(4) H(4)] N'(RK) (\Omega_4)^{-1} (\sin(\theta_4^0))^{-3} \oint_{\partial C} d^3 \eta + \right. \\
& (28) \quad \left. \frac{1}{3^2} \frac{1}{2^4} [B(4) H(4)]^2 R^2 (\Omega_5^{(C)})^{-1} \int_C d^4 \eta \right] \frac{1}{\epsilon} + F.T.
\end{aligned}$$

The infinite series over N is obviously convergent and (28) manifests, as a result, a simple pole at the limit of space-time dimensionality.

The calculational procedure for the third, fourth and fifth term in (10) is identical to that for the second. The result for the third term is

$$\begin{aligned}
& \lambda^2 B_n^4 6 \int_C d^n \eta d^n \eta' |\eta - \eta'|^{2(2-n)} \left| \frac{a_{\eta'}}{a_B} \eta - \frac{a_B}{a_{\eta'}} \eta' \right|^{2(2-n)} \rightarrow \\
& \lambda^2 \frac{1}{\pi^4} \sum_{N=0}^{\infty} \sum_{N'=0}^{N'_0} \frac{\Gamma(\frac{1}{N'_0})}{[N'^2 - N^2 + 3(N' - N)]^2} \times \\
& \left[\frac{1}{2^8} F^2(4) N'(RK^2) (\Omega_5^{(C)})^{-1} \int_C d^4 \eta + \frac{1}{2^7} [F(4) B(4) H(4)] N'(RK) (\Omega_4)^{-1} (\sin(\theta_4^0))^{-3} \oint_{\partial C} d^3 \eta + \right. \\
& (29) \quad \left. \frac{1}{3} \frac{1}{2^{10}} [B(4) H(4)]^2 R^2 (\Omega_5^{(C)})^{-1} \int_C d^4 \eta \right] \frac{1}{\epsilon} + F.T.
\end{aligned}$$

whereas the fourth and, as expected, fifth term each yield a finite result. Attention is invited at this point to the fact that the present calculation relates exclusively to the three-point function without recourse to the possibility of an overlapping divergence stemming from the self-coupling λ_0 in fig.(1d). Such a divergence would, certainly, yield an additional contribution. The exploration of such a possibility is an issue which relates directly to the renormalisation of that self-coupling and constitutes the objective of the direction in which the present technique is to be advanced. The objective of the present calculation, however, is the contribution which the zero-point function and the effective action receive exclusively due to vacuum effects at the specified loop-order. Allowing for finite contributions the stated entire contribution of the zero-point function to the bare gravitational action at third loop-order is the additive result of (25), (28) and (29). Using [12]

$$\Omega_n = \frac{2\pi^{\frac{n}{2}}}{\Gamma(\frac{n}{2})}$$

the stated contribution is

$$\begin{aligned}
I_c = & \lambda^2 \frac{1}{\pi^4} \frac{1}{2^8} \sum_{N=0}^{\infty} \left[\frac{1}{\pi^2} \frac{1}{3^3} \frac{1}{2^7} \frac{(N+2)(N+3)}{(N+1)(N+4)} (C_{N+1}^{\frac{3}{2}}(\cos\theta_4^0))^2 + \right. \\
& \sum_{N'=0}^{N'_0} \frac{1}{[N'^2 - N^2 + 3(N' - N)]^2} \left[\Gamma\left(\frac{1}{N'_0}\right) - \frac{1}{3} \frac{1}{2} \frac{\Gamma(1 + \frac{1}{N'_0})\Gamma(N' + 1 + \frac{1}{N'_0})}{\Gamma(N' + 3 + \frac{1}{N'_0})} (N+1)(N+2) \right] \times \\
& \quad 3 \frac{1}{2^3} \frac{1}{\pi^2} F^2 N' \left] \frac{RK^2}{\epsilon} \int_C d^4\eta + \right. \\
& \lambda^2 \frac{1}{\pi^6} \frac{1}{2^{13}} \sum_{N=0}^{\infty} \sum_{N'=0}^{N'_0} \frac{1}{[N'^2 - N^2 + 3(N' - N)]^2} \times \\
& \left[\Gamma\left(\frac{1}{N'_0}\right) - \frac{1}{3} \frac{1}{2} \frac{\Gamma(1 + \frac{1}{N'_0})\Gamma(N' + 1 + \frac{1}{N'_0})}{\Gamma(N' + 3 + \frac{1}{N'_0})} (N+1)(N+2) \right] (BH)^2 \frac{R^2}{\epsilon} \int_C d^4\eta + \\
& \lambda^2 \frac{1}{\pi^6} \frac{1}{2^8} \sum_{N=0}^{\infty} \sum_{N'=0}^{N'_0} \frac{1}{[N'^2 - N^2 + 3(N' - N)]^2} \times \\
& \left[\Gamma\left(\frac{1}{N'_0}\right) - \frac{1}{3} \frac{1}{2} \frac{\Gamma(1 + \frac{1}{N'_0})\Gamma(N' + 1 + \frac{1}{N'_0})}{\Gamma(N' + 3 + \frac{1}{N'_0})} (N+1)(N+2) \right] \times \\
& (FBH) N' (\sin(\theta_4^0))^{-3} \frac{RK}{\epsilon} \oint_{\partial C} d^3\eta
\end{aligned}
\tag{30}$$

with the condition $N \neq N'$ and with $n = 4$ in all n -dependent quantities.

III. Discussion

As expected [2], [3] radiative contributions on C_4 are substantially more involved than the corresponding contributions on S_4 . The complicated mathematical expression for the radiative contributions in question are the necessary consequence of both the presence of ∂C_4 and the much smaller symmetry underlying the geometry of C_4 . However, all ultra-violet divergences at third loop-order are expressed in terms of simple poles. The result expressed by (30) reveals the stated contribution of the zero-point function as the additive result of three ultra-violet divergent contributions associated with the generation of three distinct sectors in the bare and, through renormalisation, effective gravitational action. In addition to the generation of the volume divergence proportional to RK^2 there is also a volume divergence proportional to R^2 as well as a surface divergence proportional to RK . Since the one-loop contribution to the two point function, expressed by fig.(1a) and fig.(1b), is finite in the context of dimensional regularisation the associated three sectors constitute quantum corrections to the Einstein-Hilbert action generated by vacuum effects of the self-interacting scalar field at loop-orders no smaller than the third. In the context of the comparison with S_4 the volume divergence proportional to RK^2 has been exposed as the exclusive result of the presence of the boundary and, for that matter, of the cumulative effect of reflection off the boundary of signals

signifying propagation on S_4 . The presence of such a divergence in the theory is indicative of the contributions which any divergence arising from the coincidence of space-time points receives from the boundary on a general bounded manifold in addition to the contributions effected by the background geometry. With respect to the latter divergences proportional to the square of the Ricci scalar are expected on a general curved manifold [4]. However, the generation of that volume divergence on C_4 already at second order in the self-coupling λ contrasts, again, strongly with the result reported in [13] regarding the absence of any divergent contributions to the R^2 sector of the gravitational action on a general curved space-time without boundary before the fifth order in λ in the context of dimensional regularisation. The generation of a volume ultra-violet divergence substantially earlier in the perturbative expansion and, for that matter, at significantly lower energy scales is again a boundary-related effect. It becomes evident, for that matter, that the divergences inherent in the zero-point function evaluated at three-loop level receive non-trivial contributions from the boundary ∂C_4 which, effectively, dissociates the dynamical behaviour of the quantised scalar field on C_4 from that on S_4 . In the same respect the emergence of the additional surface term in the bare gravitational action as a result of the remaining surface RK -related ultra-violet divergence constitutes a purely quantum effect generated by vacuum processes in the volume of C_4 . As is the case with the RK^2 -related sector this is also a qualitatively new result effected at third loop-order by the presence of the boundary. The RK -related sector in the bare gravitational action signifies, in fact, the first surface counterterm generated by vacuum effects on a general manifold with boundary. The absence of surface counterterms on a general manifold with boundary has been reported at one-loop level [14], [15]. However, the existence of such counterterms has been anticipated at higher loop-orders on theoretical grounds [2] and the potential for their generation has been confirmed in the context of the spherical formulation for diagrammatic evaluations on a manifold with boundary [3]. The emergence of this new sector in the bare gravitational action confirms such expectations.

Two more mathematical features of the radiative contributions expressed by (30) deserve attention in their own merit. In addition to the emergence of boundary-related ultra-violet divergences the theory is characterised by the absence of infra-red divergences on C_4 . In light of (14) this result is, evidently, relevant to all orders [3]. Moreover, on the evidence of (30), all radiative contributions manifest an explicit dependence on the cut-off angular momentum for image propagation N'_0 . Such dependence is expected on a general compact bounded manifold and, necessarily, reflects the effect which the geodesic distance between points on the boundary has on the dynamical behaviour of the quantised fields.

The issue which arises as a consequence of the preceding considerations is the degree to which the results relevant to (30) can be further generalised on any curved bounded manifold. The merit of the spherical formulation which has eventuated in (30) is the expression of all Green functions of the theory in terms of the constant scalar curvature R , a result which has both formal and perturbative significance. In effect, the additional volume divergences - proportional to the Ricci and Riemann tensor-related contractions $R_{\mu\nu}R^{\mu\nu}$ and $R_{\alpha\beta\gamma\delta}R^{\alpha\beta\gamma\delta}$ respectively - which are expected on a general curved manifold are precluded on both S_4 and C_4 . However, the significance of divergences such as those

contained in (30) is, as stated, general. Their presence in this theory in the context of the demand for dimensional consistency renders them relevant to a general curved bounded manifold. Consequently, the stated K -dependent sectors are formally expected in the bare action on any curved bounded manifold and their presence in this theory is indicative of the non-trivial effect which the topology of a general bounded manifold has on the cut-off scale behaviour of quantised matter.

A relevant objection which might be raised at this point is associated with an essential aspect of the mathematical approach through which these results have been obtained. Specifically, despite its stated geometry-related generality such a conclusion would appear to still leave open the question of the regulating procedure. As stated, all ultraviolet divergences have been regulated through an analytical extension of the space-time dimensionality to an arbitrary value n . That approach preserves the maximal symmetry which underlies the spherical formulation on an arbitrary S_n and constitutes, for that matter, the natural generalisation of S_4 within the framework of dimensional regularisation. Such an extension, however, is by no means unique. The procedure of dimensional regularisation is, in fact, known to manifest a certain degree of ambiguity in curved space-time due to its inherent freedom in generalising the associated dimensionality within frameworks of different topologies [17], [18], [19]. In the present context, for example, equally valid a procedure would be the formal extension of S_4 , the covering manifold of the physical space-time C_4 , in dimensions of flat topology. Such an approach would eventuate in a regulating scheme predicated on $S_4 \times R_{n-4}$ which, obviously, does not sustain the $SO(n+1)$ de Sitter invariance manifest in (1).

The ambiguity resulting from the stated freedom of choice stems, essentially, from the effect which the presence of a non-trivial topology potentially has on the divergent structure of the theory. Such an effect is a peculiarity of the quantised theory since on the basis of the equivalence principle only local curvature terms would be expected to be of relevance to the renormalisation program. The non-trivial effect which the, global, topology potentially manifests has called into question the validity of the equivalence principle in the context of second quantisation [19], [20], [21], [22], [23]. In fact, as has already been stressed, the results obtained for the zero-point function in the context of the present calculation are indicative of such a non-trivial impact on the renormalisation program. In the context of dimensional regularisation such a topology-related impact may still persist at the limit of space-time dimensionality. In effect, both the pole structure and the resulting value for a closed-loop graph would, in general, be different for these different extensions in n dimensions. Such a difference, however, is not expected to affect the physical content of the theory. Topologically different extensions in the context of dimensional regularisation would affect, for that matter, neither the physical amplitudes manifest in the Green functions nor the renormalisation group behaviour of the theory. The perturbative evaluation of the theory's Green functions is expected to remain unaffected up to, at most, a finite renormalisation.

In view of these considerations a dimensional extension of S_4 to $S_4 \times R_{n-4}$ as opposed to S_n prior to the analytical extension of n in the context of dimensional regularisation may potentially generate a result for the three-loop contribution to the zero-point function which differs from that of (30) in pole-structure and finite part. However, the overall

contribution to the zero-point function at three-loop level will remain unaffected by these differences. It is stressed that the result in (30) represents the overall $O(\hbar^3)$ contribution of the zero-point function to the bare gravitational action. Such a contribution does not include possible other contributions to the zero-point function stemming from the scalar action. The emergence of a divergent structure different from that of (30) if $n \rightarrow 4$ is taken in the context of $S_4 \times R_{n-4}$ would be expected to cancel against the different structure which would, likewise, emerge for the overlapping divergence stemming from the bare self-interaction vertex of fig.(2d) in a manner which would eventuate in the same pole-structure for the zero-point function at three-loop level as that which would be obtained were $n \rightarrow 4$ to be taken in the context of S_n . As a result, at three-loop level the gravitational effective action on C_4 would be expected to remain unaffected by that difference in the regulating procedure up to a finite renormalisation. These remarks will be further elucidated in the context of a concrete renormalisation procedure which will constitute the natural extension of the results attained herein. Notwithstanding renormalisation, however, it might be worth conjecturing the identity of the two regulating procedures themselves in the resulting pole-structures respectively. At one-loop level the zeta-function evaluation of the effective action on a general manifold has been shown to be in agreement with the result generated by the extension of the physical dimensionality in topologically flat dimensions but not with that effected by extensions to other non-trivial topologies in the context of dimensional regularisation [17]. However, in the concrete case of the Euclidean de Sitter S_4 the zeta-function evaluation of the gravitational effective action yields a result which is in agreement with that of dimensional regularisation in the context of S_n . It would appear, in effect, that - at least at one-loop level - the two regulating procedures predicated on S_n and $S_4 \times R_{n-4}$ result in the same pole-structure. An additional argument in favour of this identity may be elicited by the absence of nonlocal, curvature-independent, divergences in the topologically non-trivial context of $S_1 \times R_3$ [23], [20], [21].

An additional ramification which the issue of the regulating procedure has relates directly to the coupling between the scalar field and the background geometry. Specifically, in conformity with dimensional regularisation, that coupling in the conformally invariant action (1) corresponds to the n -dimensional value $\xi(n) = \frac{1}{4} \frac{n-2}{n-1}$ rather than that of $\xi = \frac{1}{6}$ which relates to the improved stress tensor in four dimensions. Such an approach naturally distinguishes the three-loop contribution to the zero-point function expressed by (30) from that effected by fixing ξ to its conformal value in four dimensions. The renormalised theory, however, with its inherent one-parameter ambiguity renders the latter approach inconsistent. In fact, the immediate consequence of the theory's invariance under the renormalisation group is the scale-dependence of the theory's physical parameters. In effect, consistency with the renormalisation group invariance requires that all couplings in the theory be unspecified at the inception of the renormalisation program and, for that matter, in the semiclassical action. The extension of the coupling between the scalar field and gravity to its conformal value in n -dimensions inherent in (1) ensures that requirement in the context of dimensional regularisation while preserving the classical conformal invariance in that arbitrary dimensionality. Radiative effects can, of course, still be computed with $\xi = \frac{1}{6}$ from the outset. Such an approach, however,

trivialises the renormalisation group behaviour of the theory unless that classical value for ξ also corresponds to a fixed point for the renormalised theory, a situation which amounts, essentially, to a tautology considering that the evaluation of any possible fixed point necessitates the solution of the renormalisation group equations. Again, these remarks will be further elucidated in the context of a concrete calculation which should render the renormalisation group behaviour of the theory manifest.

Pursuant to the question of generality it is also worth mentioning that results in perturbative renormalisation relevant to a massless scalar field conformally coupled to the background geometry of C_4 at semi-classical level with a ϕ^4 self-interaction are of immediate importance to quantum cosmology in view of the description which the ground state of the cosmological wavefunction admits in terms of constant-curvature gravitational instantons in imaginary time. The results hitherto attained on C_4 up to third loop-order and the stated degree of generality which they, qualitatively, admit on any curved bounded manifold call into question the issue of renormalisation at that loop-order. It is evident from (30) that the ultra-violet divergences inherent in the zero-point function necessitate, at that order, the simultaneous introduction of both volume and surface counterterms. This result indicates that, in conformity with theoretical expectation [2], [3], renormalisation in the volume and surface effective action is accomplished simultaneously. Such a situation shall be explicitly confirmed in the context of a concrete renormalisation program which, as stated, shall also advance the remarks made above as to the relation between the present results and the regulating procedure. The task of renormalising the theory past one-loop level is rather arduous in view of the complicated associated expressions such as that in (30). However, work on these lines is in progress.

REFERENCES

- [1] I. G. Avramidi and G. Esposito, *Class. Quant. Gravity* **15** 281, (1998)
- [2] G. Tsoupros, *Class. Quantum Gravity* **17** 2255-2266, (2000)
- [3] G. Tsoupros, e-print Archive: hep-th/0107019
- [4] See for example: N. D. Birrel and P. C. W. Davies, **Quantum Fields in Curved Space** Cambridge Monographs in Mathematical Physics (1984) and references therein.
- [5] D. G. C. McKeon and G. Tsoupros, *Class. Quantum Gravity* **11** 73-81, (1994)
- [6] I. Drummond, *Nucl. Phys. B* **94** 115, (1975)
- [7] I. Drummond, *Phys. Rev. D* **19**, 4 1123, (1979)
- [8] I. T. Drummond and G. M. Shore, *Phys. Rev. D* **19**, 4 1134, (1979)
- [9] G. M. Shore, *Ann. Phys. (NY)*, **117** 89, (1979)
- [10] G. M. Shore, *Phys. Rev. D* **21** 2226 (1980)
- [11] G. 't Hooft *Nucl. Phys. B* **61** 455, 1973
- [12] See for example the **Higher Transcendental Functions** *The Bateman Manuscript Project*, (ed. A. Erdelyi), McGraw Hill, New York, (1953), vol. II; also N. J. Vilenkin **Special Functions and the Theory of Group Representations** The American Mathematical Society (1968)
- [13] L. S. Brown and J. C. Collins *Ann. Phys.* **130**, 215-248 (1980)
- [14] S. D. Odintsov, *Class. Quant. Gravity* **7** 445-448 (1990)
- [15] S. D. Odintsov and A. Wipf, *Phys. Lett. B* **356** 26-31 (1995)
- [16] A. O. Barvinsky and S. N. Solodukhin, *Nucl. Phys. B* **479** 305-318, (1996) (e-Print Archive: gr-qc/9512047)
- [17] S. W. Hawking, *Commun. math. Phys.* **55**, 133-148 (1977)
- [18] J. S. Dowker and R. Critchley, *Phys. Rev. D* **13**, 12, 3224-3232 (1975)
- [19] D. J. Toms, *Phys. Rev. D* **26**, 10, 2713-2729 (1982)

- [20] D. J. Toms, Phys. Rev.D **21**, 928 (1980)
- [21] D. J. Toms, Ann. Phys. (N.Y.)**129**, 334 (1980)
- [22] I. T. Drummond and S. J. Hathrell, Phys. Rev. D **22**, 343 (1980)
- [23] N. D. Birrell and L. H. Ford, Phys. Rev. D **22**, 330 (1980)